

Hox Genes, Digit Identities and the Theropod/Bird Transition

FRIETSON GALIS^{1*}, MARTIN KUNDRÁT² AND JOHAN A.J. METZ^{1,3}

¹*Institute of Biology, Leiden University, 2300 RA Leiden, The Netherlands*

²*Department of Zoology, Natural Science Faculty, Charles University, Prague, Czech Republic*

³*Adaptive Dynamics Network, Institute for Applied Systems Analysis, A-2361 Laxenburg, Austria*

ABSTRACT Vargas and Fallon (2005. *J Exp Zool (Mol Dev Evol)* 304B:86–90) propose that *Hox* gene expression patterns indicate that the most anterior digit in bird wings is homologous to digit 1 rather than to digit 2 in other amniotes. This interpretation is based on the presence of *Hoxd13* expression in combination with the absence of *Hoxd12* expression in the second digit condensation from which this digit develops (the first condensation is transiently present). This is a pattern that is similar to that in the developing digit 1 of the chicken foot and the mouse hand and foot. They have tested this new hypothesis by analysing *Hoxd12* and *Hoxd13* expression patterns in two polydactylous chicken mutants, *Silkie* and *talpid2*. They conclude that the data support the notion that the most anterior remaining digit of the bird wing is homologous to digit 1 in other amniotes either in a standard phylogenetic sense, or alternatively in a (limited) developmental sense in agreement with the Frameshift Hypothesis of Wagner and Gautier (1999, i.e., that the developmental pathway is homologous to the one that leads to a digit 1 identity in other amniotes, although it occurs in the second instead of the first digit condensation). We argue that the *Hoxd12* and *Hoxd13* expression patterns found for these and other limb mutants do not allow distinguishing between the hypothesis of Vargas and Fallon (2005. *J Exp Zool (Mol Dev Evol)* 304B:86–90) and the alternative one, i.e., the most anterior digit in bird wings is homologous to digit 2 in other amniotes, in a phylogenetic or developmental sense. Therefore, at the moment the data on limb mutants does not present a challenge to the hypothesis, based on other developmental data (Holmgren, 1955. *Acta Zool* 36:243–328; Hinchliffe, 1984. In: Hecht M, Ostrom JH, Viohl G, Wellnhofer P, editors. *The beginnings of birds*. Eichstätt: Freunde des Jura-Museum. p 141–147; Burke and Feduccia, 1997. *Science* 278:666–668; Kundrát et al., 2002. *J Exp Zool (Mol Dev Evol)* 294B:151–159; Larsson and Wagner, 2002. *J Exp Zool (Mol Dev Evol)* 294B:146–151; Feduccia and Nowicki, 2002. *Naturwissenschaften* 89:391–393), that the digits of bird wings are homologous to digits 2,3,4 in amniotes. We recommend further testing of the hypothesis by comparing *Hoxd* expression patterns in different taxa. *J. Exp. Zool. (Mol. Dev. Evol.)* 304B:198–205, 2005. © 2005 Wiley-Liss, Inc.

Vargas and Fallon (2005) propose a new significance to the expression of *Hoxd12* and *Hoxd13* gene expression patterns in the developing digits of forelimbs and hindlimbs of amniotes. They propose that the combined expression pattern of these genes provides a reliable marker either of the condensation homologous in a phylogenetic sense to that from which the most anterior digit develops in pentadactyl amniotes (i.e., condensation 1 of digit 1), or of the condensation from which a digit develops with a digit 1 identity, i.e., with the same adult identity as that of digit 1 (in a phylogenetic sense), regardless of the condensation from which the digit develops (see our

definition of concepts in the appendix). The second proposal by Vargas and Fallon (2005) is difficult to evaluate, because of the evolutionary diversity of digit 1 identities among amniotes, which is manifested in a large variation of shapes, sizes and phalanx numbers. However, the proposal may be evaluated in taxa in which there has been relatively little evolutionary change in forelimbs and/or hindlimbs. In practice, we follow Tabin ('92)

*Correspondence to: Frietson Galis, Institute of Biology, Leiden University, P.O. Box 9516, 2300 RA Leiden, The Netherlands.

E-mail: galis@rulsfb.leidenuniv.nl

Received 3 January 2005; Accepted 14 February 2005

Published online 6 May 2005 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/jez.b.21042.

and Litingtung et al. (2002) in that the identity of a digit is inferred from the number of phalanges, the size, shape and structure of the adult digit.¹

Hox genes and digit identity

During limb development in mice and chickens, 5' group *Hox* genes play an important role in the antero-posterior (A-P) patterning of the limbs, including the digits. Recently, much progress has been made in understanding the determination of digit identity and several new models have been proposed (e.g., Sanz-Ezquerro and Tickle, 2003; Suzuki et al., 2004; Harfe et al., 2004; Ahn and Joyner, 2004; Zakany et al., 2004). The full picture of digit identity determination and in particular the role of *Hox* genes is, however, unclear. Unravelling the role of *Hox* genes in the A-P patterning of digits is difficult, because the signalling of these genes forms part of a complicated and dynamically changing feedback-loop that involves many other A-P patterning genes, amongst others *Shh* and *Bmp* genes (e.g., Drossopoulou et al., 2000; Chiang et al., 2001; Litingtung et al., 2002; Te Welscher et al., 2002; Ros et al., 2003; Ahn and Joyner, 2004; Harfe et al., 2004; Selever et al., 2004; Tickle, 2004; Zakany and Duboule, 1999; Zakany et al., 2004). The complexity of the interactions is further increased by the integration of A-P patterning with dorsoventral and proximodistal patterning (Tickle, 2004) and by the high functional redundancy of *Hox* genes (Fig. 1, Davis and Capecchi, 1996; Goff and Tabin, 1997; Zakany and Duboule, 1999; Kmita et al., 2002). Only if the *Hox* genes are the first genes to be expressed in a clear, differentiated manner correlated with digit identity, can they tentatively be inferred to steer the local network leading to the digit form. As long as data of the required fine temporal resolution are lacking, the *Hox* expression pattern can only be viewed as a useful correlate of digit identity.

Vargas and Fallon (2005) draw attention to interesting similarities of *Hoxd12* and *Hoxd13* expression patterns in the developing digits of forelimbs and hindlimbs of mice and chickens. *Hoxd13* is present in all developing digit condensations whereas *Hoxd12* has a more posterior boundary and is not expressed in the most anterior condensation of the remaining digits, i.e., the second digit condensation in the chicken




Fig. 1. Dorsal view of the forelimbs of a wild-type mouse and of three mutant mouse strains produced by gene targeting (*Hoxd11*^{-/-}, *Hoxd12*^{-/-} and *Hoxd13*^{-/-}). Loss of *Hoxd11* or *Hoxd12* in the mutants hardly affects the phenotype of the digits, suggesting functional redundancy among posterior *Hox* genes. The phenotype of the limb of *Hoxd13*^{-/-} mice is more disturbed (from Davis and Capecchi, '96).

wing (the first condensation is transiently present and disappears at an early stage; Prein, '14; Holmgren, '55; Feduccia and Nowicki, 2002; Kundrát et al., 2002; Larsson and Wagner, 2002) and the first digit condensation in the hand and foot of the mouse and the foot of the chicken.

Proposal of two scenarios

Vargas and Fallon now propose in their first scenario that in birds and mammals the absence of *Hoxd12* expression combined with the presence of *Hoxd13* expression in this condensation should not only be seen as a marker of the most anterior condensation of the remaining digits, but specifically of the condensation of digit 1 (in a phylogenetic sense), i.e., the local *Hox* expression pattern indicates the condensation of digit 1 in amniotes, regardless of the differences in identity of the adult digit 1 between taxa and between forelimbs and hindlimbs. They deduce from this that the second condensation in the developing bird would then be the condensation of digit 1 and the first condensation that of the prepollex. In this scenario, bird wings would not have digits 2, 3, 4, as commonly assumed by developmental biologists (e.g., Burke and Feduccia, '97; Hinchliffe, '97; Wagner and Gauthier, '99; Kundrát et al., 2002), but digits 1, 2, 3 (although 1, 3, 4 and 1, 2, 4 are also possible). This interpretation would remove the long-standing problem of how birds, presumed to have digits 2, 3, 4 remaining at the end of evolution, can have descended from theropods that are generally assumed to have had digits 1, 2, 3 (e.g., Wagner and Gauthier, '99; Zhou, 2004). This problem stems from the difference between a homology interpretation of digits in birds that is

¹Position is also a form criterion, but the antero-posterior position of the digit is excluded here.

based on a comparison of early amniote development, in particular, of the branching pattern and the number of the digit condensations from which condensations 2, 3, 4 continue developmentally, and one that is based on a comparison of the adult morphology of the digits of birds and theropods, where in theropods and birds adult digits 1, 2, 3 appear to have remained in the fossil record and digits 4 and 5 are reduced (Wagner and Gauthier, '99).

In an alternative scenario, Vargas and Fallon (2005) propose that the mutant data also supports the Frameshift Hypothesis by Wagner and Gauthier ('99). This hypothesis states that the digits of bird wings have undergone homeotic identity shifts in their evolutionary past such that a digit with the adult identity of digit 1 develops in the condensation formerly of digit 2, a digit with the adult identity of digit 2 in the condensation formerly of digit 3 and a digit with the adult digit 3 identity in the condensation formerly of digit 4. This hypothesis was specifically proposed to solve the above-mentioned discrepancy between the homology interpretations based on developmental data of amniotes (2, 3, 4) and paleontological data of adult theropods and birds (1, 2, 3). For this alternative scenario, Vargas and Fallon (2005) thus hypothesize that in amniotes the absence of *Hoxd12* expression combined with the presence of *Hoxd13* expression indicates in forelimbs and hindlimbs the condensation from which a digit develops with the identity of a digit 1, rather than that of digits 2, 3, 4 or 5, regardless of the position of the digit condensation. They have tested their hypothesis by analysing *Hoxd12* and *Hoxd13* expression patterns in two polydactylous chicken mutants, Silkie and *talpid2*.

ANALYSIS OF CHICKEN AND MICE MUTANTS

Does data on polydactylous chickens support the hypothesis?

Silkie

In Silkie chickens, the most common variant has feet with an extra digit, with three phalanges anterior to digit 1. The extra digit is morphologically very similar to digits 2, 3 and 4 of the ordinary foot. Most authors interpret the Silkie polydactyly as a partial mirror image duplication of the foot with a mirror image digit 2 anterior to digit 1 (i.e., digits with identities 2, 1, 2, 3, 4; Grönberg, 1894; Braus, '08; Gabriel, '46; Land-

auer, '48). They base this on the range of digit variations displayed by both homozygous Silkie chickens and hybrids with normal-toed breeds. Indeed, the *Hoxd12* and *Hoxd13* patterns in Silkie are mirrored in the duplicated digit 2 condensation, i.e., both *Hoxd12* and *Hoxd13* are expressed. In experimentally induced full and partial mirror image duplications of chick wings, *Hoxd11*, *Hoxd12* and *Hoxd13* patterns are similarly mirrored during development (Izpisúa Belmonte et al., '91; Nohno et al., '91). If indeed the polydactyly in the Silkie mutant should be interpreted as a partial mirror image duplication, the *Hoxd12* and *Hoxd13* patterns in the developing extra digit do not seem to provide support for Vargas and Fallon's hypothesis, because the mirror image duplication is most likely induced upstream of the *Hox* gene expression at embryonic day 8 (e.g., by ectopic anterior *Shh* expression), allowing the possibility that digit identity is induced by the same upstream stimuli, at least in part independent of the *Hox* expression. It is thus necessary to investigate first whether or not the Silkie polydactyly should be seen as a mirror image duplication and what induces the mirror duplication, before the Silkie data can be adduced as an argument in favour of Vargas and Fallon's hypothesis.

Talpid2

In chicken *talpid2* mutants, the limbs have many digits that have lost wild-type morphological identity and that are fused (Fig. 2; Goetinck and Abbott, '64; MacCabe and Abbott, '74; Dvorak and



Fig. 2. From left to right, *talpid2* wing, wild-type wing, *talpid2* leg, wild-type leg. *Talpid2* limbs have supernumerary digits that have lost wild-type morphological identity (from Goetinck and Abbott, '64).

Fallon, '91; Caruccio et al., '99). The A-P polarity is disturbed and weakened, which is assumed to be causally implicated in the loss of wild-type identity. In the foot the digits mostly consist of three phalanges, although some phalanges (and some digits) disappear during development due to apoptosis. Vargas and Fallon assume that the first digit has become posteriorized in *talpid2* mutant feet, because in all digits the number of phalanges is three as in wild-type posterior digits. *Hoxd12* and *Hoxd13* are expressed in all digits on day 7 and this is in agreement with Vargas and Fallon's hypothesis, if indeed the assumption holds true that the first digit has assumed the identity of more posterior digits. We feel that the number of phalanges is not sufficient to conclude this, given the loss of wild-type identity of all digits, including the most anterior digit. The assumption of a posteriorization of the identity of the first digit is not supported by the majority of anatomical characters that determine the identity of a digit (Fig. 2).

Phalanx number, digit identity and Hox genes

Regarding the relationship between phalanx number and digit identity, it is important to note that in chicken wings the number of phalanges of adult digits has been reduced compared with *Archaeopteryx* (phalangeal formulas resp. 1(2), 2, 1 and 2, 3, 4). Nonetheless, the evolutionary reduction that took place in the middle digit of birds from three phalanges to two has never been considered sufficient reason to conclude that the identity of the digit has been changed into that of the most anterior digit, because of the differences between other anatomical characters. In addition, the *Hoxd12* and *Hoxd13* expression patterns do not support this, because in the chicken wing both *Hoxd12* and *Hoxd13* are expressed in the condensation of the second and third remaining digits. Hence, in chicken limbs, the combination of both *Hoxd12* and *Hoxd13* expression is associated with the development of digits with one, two and three phalanges. Therefore, there is no tight correlation between phalanx number and *Hoxd12* and *Hoxd13* gene expression in chicken limbs.

Analysis of other chicken limb mutants

Analysis of other chicken and mouse limb mutants provides evidence that a digit 1 identity (including a phalanx number of two) can apparently develop in the presence of *Hoxd12* expres-

sion and, in addition, that the identity of digits 2, 3 and 4 (including a phalanx number of three) can develop in the absence of *Hoxd12* expression. Therefore, in limb mutants in general, there does not appear to be a very tight correlation between the *Hoxd12* and *Hoxd13* gene expression and digit identity (including phalanx number). Suzuki et al. (2004) show, for example, that misexpression of *Tbx2* leads to an anterior expansion of *Hoxd12* expression without a posteriorization of the identity of digit 1 (mutants have feet with digits with identities 1, 3, 4, 4 and 1, 2, 4, 4). Selever et al. (2004) show that in mutant limbs in which *Bmp* expression is inactivated (*Prx1cre; Bmp4n/f* mutants), complete and partial posteriorizations of the identity of digit 1 occur in the absence of a change in the expression pattern of *Hoxd12* and *Hoxd13* gene expression. Similarly, it has been shown that over-expression of *Hoxd11* leads to partial or complete posteriorization of the identity of digit 1 (Morgan et al., '92; Goff and Tabin, '97), presumably in the absence of any changes in *Hoxd12* and *Hoxd13* expression (Morgan et al., '92). Finally, Davis and Capecchi ('96) show that in mutants in which *Hoxd12* expression is entirely absent (*Hoxd12*^{-/-}), there is virtually no change in the identity of the digits (Fig. 1). The results emphasize the functional redundancy of posterior *Hoxd* genes in the limb. This redundancy also follows from the fact that phenotypic effects in the digits are again very limited when *Hoxd13* is absent and *Hoxd12* expression is expanded anteriorly, similarly to the normal expression of *Hoxd13* (as a result of the deletion of the *Hoxd13* locus; Kmita et al., 2002). This again limits the reliability of the *Hoxd12* and *Hoxd13* patterns as sole indicators of digit identity in limb mutants.

The analyses of these and other mutant limbs show that *Hoxd12* and *Hoxd13* play an important role in the patterning of the A-P identity of the digits. However, they do not allow the conclusion that the presence of *Hoxd13* expression together with the absence of *Hoxd12* in mice and chicken mutant digit condensations provides molecular evidence for the development of a digit 1 identity, to the exclusion of a more posterior digit identity.

FIVE INITIAL CONDENSATIONS IN THE BIRD WING

Prepollex in bird wings?

The ossifying digits in the bird wings develop, as in the feet, in the middle three of the initial five digital mesenchymal condensations (e.g., Prein, '14;




Fig. 3. Comparison of the wing (top row) and foot (bottom row) development in the chicken, dorsal views. Digit 4 is labelled in each specimen. Note the strong resemblance of the patterns. In the foot, the transient condensation of digit 5 can be seen. (Reprinted with permission from Burke AC, Feduccia A. 1997. Developmental patterns and the identification of homologies in the avian hand. *Science* 278:666–668). Copyright 1997 AAAS.

Holmgren, '55; Feduccia and Nowicki, 2002; Kundrát et al., 2002; Larsson and Wagner, 2002). Vargas and Fallon (2005) suggest that the most anterior condensation, which is only transiently present, may represent the prepollex rather than the digit 1 condensation as is generally assumed. This is highly improbable, because this would imply that digit condensations are present of the prepollex (digit 0) and digits 1, 2, 3 and 4 and not of digit 5. The general consensus is that the prepollex was lost much earlier than digit 5 in the ancestral lineage of birds and reptiles (e.g., Romer, '55; Wagner and Gauthier, '99). Moreover, evolutionary digit reduction tends to be extremely slow and gradual (Lande, '78; Galis et al., 2002). Together, these two facts suggest that there should still be a condensation present of the most recently lost digit 5 rather than one of the prepollex. The interpretation of a prepollex condensation is also not in agreement with the similarity of the branching pattern of the digit condensations in the chicken foot and wing and in other amniote limbs that also indicate that the remaining digits in the bird wings develop from digit condensations 2, 3 and 4 (Fig. 3; Hinchliffe, '84; Burke and Feduccia, '97; Kundrát et al., 2002). This supports the notion that bird digits are digits 2, 3, 4 in a phylogenetic homological sense.

DISCUSSION

On the basis of *Hoxd12* and *Hoxd13* expression patterns, the data on limb mutants unfortunately

do not allow distinguishing between the hypothesis that the most anterior digit in the wing of birds is homologous to digit 1 in other amniotes, and the alternative that it is homologous to digit 2, either in a phylogenetic or a developmental sense. More particularly, the disturbed nature of the digital patterning in most discussed mutants makes a proper evaluation of Vargas and Fallon's hypothesis difficult. A more straightforward test of the hypothesis would be to investigate *Hoxd12* and *Hoxd13* gene expression patterns in amniote species other than chickens and mice. In particular, it would be useful to examine species from taxa that have independently lost digit 1 in forelimbs or hindlimbs during evolution (e.g., ostriches, dogs and many reptilian species). Furthermore, it would be useful to also include analysis of *Hox* gene expression patterns (including *Hoxd11*) at earlier stages, when much of the digit identity has not yet been specified (e.g., the number of phalanges, Hartmann and Tabin, 2001; see also Dudley et al. (2002) and Zakany et al. (2004) for early specification), even though it has been shown that at later stages part of the specification has not yet irreversibly happened (Dahn and Fallon, 2000).

Unfortunately, the late *Hox* activity cannot be causally implicated in the initial specification of the digit identity anymore, although it is of course possible that it is correlated with it.

At the moment the data on mutant limbs does not present a challenge to the hypothesis that is based on developmental data; that is, the digits of

bird wings are homologous to digits 2, 3, 4 in other amniotes. This appears to leave the problem of how birds with digits 2, 3, 4 in their forelimbs can have descended from theropods that are generally assumed to have digits 1, 2, 3. However, the apparent conflict between homology based on developmental and paleontological data does not mean that we should seriously question the descent of birds from theropods. In an earlier paper (Galis et al., 2003), we have discussed two scenarios that reconcile a theropod descent of birds with the digit identities that have been found: (a) the above-mentioned Frameshift Hypothesis by Wagner and Gauthier ('99) and (b) the hypothesis that the theropod ancestors of birds may have had hands with digits 2, 3, 4, rather than 1, 2, 3.

Frameshift Hypothesis

The Frameshift Hypothesis has several problems. The homeotic identity shifts cannot be achieved by straightforward identity shifts in four consecutive digits alone. The frameshift also requires (a) the reversal of the evolutionary reduction of digit 4 into a fully functional digit (a polydactylous change) and (b) the induction of digit reduction in digit 1 (an oligodactylous change). Both polydactylous and oligodactylous changes are highly constrained as single mutational steps and have never been documented in amniotes at a species level, despite the common occurrence of such mutations within species (i.e., high intraspecific and no interspecific variation, Lande, '78; Galis et al., 2001, 2002). In addition, a homeotic shift of digits 1, 2, 3 into digits 2, 3, 4 in theropods without further anatomical changes does not appear to lead to an adaptive advantage. Therefore, it appears probable that not one, but several highly constrained mutational changes are necessary to achieve the proposed frameshift, without indications, as yet, for a selective advantage that would favour these changes.

Theropod ancestors with wing digits 2, 3, 4

The scenario that theropod ancestors of birds may have had hands with digits 2, 3, 4 is also problematic. Analysis of the fossil data strongly suggests the reduction of digits 4 and 5 in pentadactyl theropods. However, the reduction of digit 4 is not absolutely certain, and, given the gaps in the Jurassic fossil record of theropods, a bilateral reduction of digits 1 and 5 cannot be entirely excluded (Galis et al., 2003). Given the plausibility of a descent of birds from theropods

and the arguably low mechanistic plausibility of the Frameshift Hypothesis, we think that the hypothesis that theropod ancestors of birds had hands with digits 2, 3, 4 should be explored more thoroughly (see also Zhou, 2004).

Plausibility of evolutionary transitions

Larsson and Wagner (2003) in their reply to Galis et al. (2003) disagree with this reasoning and suggest that the inferred low mechanistic plausibility of the frameshift hypothesis is less relevant as it may be inaccurate due to the incompleteness of our scientific knowledge. Our opinion is that considerations about the plausibility of evolutionary transitions on morphogenetic or selective grounds should be an essential ingredient of any attempt to reconstruct such transitions in a phylogenetic context. In this context, it may be noted that the Frameshift Hypothesis was itself also put forward by Wagner and Gauthier ('99) to make the evolutionary transition from theropod hands with digits 1, 2, 3 to bird wings with digits 2, 3, 4 mechanistically plausible. Not only that, consideration of the plausibility of evolutionary transitions is actually a routine procedure (though generally unacknowledged as such) in the construction of cladograms when choices are made between the multiple loss or multiple gain of complex traits.

Both scenarios, thus, currently lack convincing support. Hopefully new paleontological data may provide either a plausible adaptive scenario for the frameshift, or support for a bilateral reduction in the hands of the theropod ancestors of birds (digits 2, 3, 4) or for yet another hypothesis. In addition, further testing of Vargas and Fallon's hypothesis in amniote species with independently reduced numbers of digits will provide new information on the molecular basis of evolutionary digit reduction, and hence, on the transition of theropod digits to bird digits. For the moment, the enigma persists.

ACKNOWLEDGMENTS

We thank Annie Burke, Jacques van Alphen and Elizabeth van Ast for useful comments on the manuscript, and Linda Oswald for kindly providing a scan of Fig. 1.

LITERATURE CITED

Ahn S, Joyner AL. 2004. Dynamic changes in the response of cells to positive hedgehog signaling during mouse limb patterning. *Cell* 118:505–516.

- Braus H. 1908. Entwicklungsgeschichtliche analyse der hyperdaktylie. Muenchener Medizinische Wochenschrift 55:386–390.
- Burke AC, Feduccia A. 1997. Developmental patterns and the identification of homologies in the avian hand. *Science* 278:666–668.
- Caruccio NC, Martinez-Lopez A, Harris M, Dvorak L, Bitgood J, Simandl BK, Fallon JF. 1999. Constitutive activation of sonic hedgehog signaling in the chicken mutant talpid(2): Shh independent outgrowth and polarizing activity. *Dev Biol*. 212:137–149.
- Chiang C, Litingtung Y, Harris MP, Simandl BK, Li Y, Beachy PA, Fallon JF. 2001. Manifestation of the limb prepatterning: limb development in the absence of sonic hedgehog function. *Dev Biol* 236:421–435.
- Dahn RD, Fallon JF. 2000. Interdigital regulation of digit identity and homeotic transformation by modulated BMP signaling. *Science* 289:438–441.
- Davis AP, Capecchi MR. 1996. A mutational analysis of the 5' HoxD genes: dissection of genetic interactions during limb development in the mouse. *Development* 122:1175–1185.
- Drossopoulou G, Lewis KE, Sanz-Ezquerro JJ, Nikbakht N, McMahon AP, Hofmann C, Tickle C. 2000. A model for anteroposterior patterning of the vertebrate limb based on sequential long- and short-range Shh signalling and Bmp signalling. *Development* 127:1337–1348.
- Dudley AT, Ros MA, Tabin CJ. 2002. A re-examination of proximodistal patterning during vertebrate limb development. *Nature* 418:539–544.
- Dvorak L, Fallon JF. 1991. Talpid2 mutant chick limb has anteroposterior polarity and altered patterns of programmed cell death. *Anat Rec* 231:251–260.
- Feduccia A, Nowicki J. 2002. The hand of birds revealed by early ostrich embryos. *Naturwissenschaften* 89:391–393.
- Gabriel ML. 1946. The effect of local applications of colchicine on Leghorn polydactylous chick embryos. *J Exp Zool* 101:339–350.
- Galis F, van Alphen JJM, Metz JAJ. 2001. Why five fingers? Evolutionary constraints on digit numbers. *Trends Ecol Evol* 16:637–646.
- Galis F, van Alphen JJM, Metz JAJ. 2002. Digit reduction: via repatterning or developmental arrest. *Evol Dev* 4, 249–251.
- Galis F, Kunderát, M, Sinervo B. 2003. An old controversy solved: bird embryos have five fingers. *Trends Ecol Evol* 18:7–9.
- Goetinck PF, Abbott, UK. 1964. Studies on limb morphogenesis. I. Experiments with the polydactylous mutant, talpid. *J Exp Zool* 155:161–170.
- Goff DJ, Tabin CJ. 1997. Analysis of Hoxd-13 and Hoxd-11 misexpression in chick limb buds reveals that Hox genes affect both bone condensation and growth. *Development* 124:627–636.
- Grönberg G. 1894. Beiträge zur kenntniss der polydactylen hünerrassen. *Anat Anz* 9:509–516.
- Harfe BD, Scherz PJ, Nissim S, Tian H, McMahon AP, Tabin CJ. 2004. Evidence for an expansion-based temporal Shh gradient in specifying vertebrate digit identities. *Cell* 118:517–528.
- Hartmann C, Tabin CJ. 2001. Wnt-14 plays a pivotal role in inducing synovial joint formation in the developing appendicular skeleton. *Cell* 104:341–351.
- Hinchliffe JR. 1984. “One, two, three” or “Two, three, four”: an embryologist’s view of the homologies of the digits and carpus of modern birds. In: Hecht M, Ostrom JH, Viohl G, Wellenhofer P, editors. *The beginnings of birds*. Eichstätt: Freunde des Jura-Museum. p 141–147.
- Hinchliffe JR. 1997. The forward march of the bird-dinosaurs halted? *Science* 278:596–597.
- Holmgren N. 1955. Studies on the phylogeny of birds. *Acta Zool* 36:243–328.
- Izpisúa Belmonte JC, Tickle C, Dollé P, Wolpert L, Duboule D. 1991. Expression of the homeobox Hox-4 genes and the specification of position in chick wing development. *Nature* 350:585–589.
- Kmita M, Fraudeau N, Herault Y, Duboule D. 2002. Serial deletions and duplications suggest a mechanism for the collinearity of Hoxd genes in limbs. *Nature* 420:145–150.
- Kunderát M, Seichert V, Russell AP, Smetana K, Jr. 2002. Pentadactyl pattern of the avian wing autopodium and pyramid reduction hypothesis. *J Exp Zool (Mol Dev Evol)* 294B:151–159.
- Landauer W. 1948. The phenotypic modification of hereditary polydactylism of fowl by selection and insulin. *Genetics* 33:133–157.
- Lande R. 1978. Evolutionary mechanisms of limb loss in tetrapods. *Evolution* 32:73–92.
- Larsson HCE, Wagner GP. 2002. Pentadactyl pattern of the avian wing autopodium and pyramid reduction hypothesis. *J Exp Zool (Mol Dev Evol)* 294B:146–151.
- Larsson HCE, Wagner GP. 2003. Old morphologies misinterpreted. *Trends Ecol Evol* 18:10.
- Litingtung Y, Dahn RD, Li Y, Fallon JF, Chiang C. 2002. Shh and Gli3 are dispensable for limb skeleton formation but regulate digit number and identity. *Nature* 418:979–983.
- MacCabe JA, Abbott UK. 1974. Polarizing and maintenance activities in two polydactylous mutants of the fowl: diplopodia and talpid. *J Embryol Exp Morphol* 31:735–746.
- Morgan BA, Izpisúa-Belmonte J-C, Duboule, D, Tabin CJ. 1992. Targeted misexpression of *Hox-4.6* in the avian limb bud causes apparent homeotic transformations. *Nature* 358:236–239.
- Nohno T, Noji S, Koyama E, Ohyama K, Myokai F, Kuroiwa A, Saito T, Taniguchi S. 1991. Involvement of the *Chox-4* chicken homeobox genes in determination of anteroposterior axial polarity during limb development. *Cell* 64: 1197–1205.
- Prein, F. 1914. Die entwicklung des vorderen extremitaetenskelettes beim haushuhn. In: Merkel F, Bonnet R, editors. *Anatomische hefte. Beitrage und referate zur anatomie und entwicklungsgeschichte*. Wiesbaden: J.F. Bergmann.
- Romer AS. 1955. *Osteology of the reptiles*. Chicago: University of Chicago Press.
- Ros MA, Dahn RD, Fernandez-Teran M, Rashka K, Caruccio NC, Hasso SM, Bitgood JJ, Lanman JJ, Fallon JF. 2003. The chick oligozeugodactyly (ozd) mutant lacks sonic hedgehog function in the limb. *Development*. 130:527–537.
- Sanz-Ezquerro JJ, Tickle C. 2003. Fgf signaling controls the number of phalanges and tip formation in developing digits. *Curr Biol* 13:1830–1836.
- Selever J, Liu W, Lu M-F, Behringer RR, Martin JF. 2004. *Bmp4* in limb bud mesoderm regulates digit pattern by controlling AER development. *Dev Biol* 276:1–12.
- Suzuki T, Takeuchi J, Koshiba-Takeuchi K, Ogura T. 2004. *Tbx* genes specify posterior digit identity through Shh and BMP signaling. *Dev Cell* 6:43–53.
- Tabin CJ. 1992. Why we have (only) five fingers per hand: hox genes and the evolution of paired limbs. *Development* 116:289–296.

- Tickle C. 2004. The contribution of chicken embryology to the understanding of vertebrate limb development. *Mech Dev* 121:1019–1029.
- Vargas AO, Fallon JF. 2005. Birds have dinosaur wings: the molecular evidence. *J Exp Zool (Mol Dev Evol)* 304B:86–90.
- Wagner GP, Gauthier JA. 1999. 1,2,3=2,3,4: a solution to the problem of the homology of the digits in the avian hand. *Proc Natl Acad Sci USA* 96:5111–5116.
- Te Welscher P, Zuniga A, Kuijper S, Drenth T, Goedemans HJ, Meijlink F, Zeller R. 2002. Progression of vertebrate limb development through SHH-mediated counteraction of GLI3. *Science* 298:827–830.
- Zakany J, Duboule D. 1999. Hox genes in digit development and evolution. *Cell Tissue Res* 296:19–25.
- Zakany J, Kmita M, Duboule D. 2004. A dual role for Hox genes in limb anterior-posterior asymmetry. *Science* 304:1669–1672.
- Zhou Z. 2004. The origin and early evolution of birds: discoveries, disputes, and perspectives from fossil evidence. *Naturwissenschaften* 91:455–471.

APPENDIX

CONCEPTS

There are four concepts that we have to deal with, condensation number and identity and digit number and identity. Digit refers to the adult, condensation to the earliest recognizable shape in the embryo. Numbers are observationally determined, ordinal numbers counted antero-posteriorly. Identities are determined by phylogenetic homology with adult mice as a reference. Condensations in mice derive their identity from the corresponding adult. Two features, be they condensations or adult digits, in a parent and offspring

are called homologous if they are closely similar. Two features in arbitrary organisms are called homologous if they can be linked through a phylogenetic chain of parent–offspring pairs. In the case of condensations, similarity is determined by the combination of the four form parameters (position, size, shape and structure). In the case of digits, similarity is based on adult form parameters only. (The implicit assumption is thus that over the generations the disappearance of condensations and the change of adult shapes occurs only gradually.) Disappearance of the prepollex leads to a shift of digit number relative to digit identity. A shift of digit identity relative to that of the corresponding condensation is called a frameshift.