

Do Large Dogs Die Young?

FRIETSON GALIS^{1*}, INKE VAN DER SLUIJS¹, TOM J.M. VAN DOOREN¹, JOHAN A.J. METZ^{1,2}, AND MARC NUSSBAUMER³

¹*Institute of Biology, Leiden University, 2300 RA Leiden, The Netherlands*

²*Adaptive Dynamics Network, Institute for Applied Systems Analysis, A-2361 Laxenburg, Austria*

³*Natural History Museum, CH-3005 Berne, Switzerland*

ABSTRACT In most animal taxa, longevity increases with body size across species, as predicted by the oxidative stress theory of aging. In contrast, in within-species comparisons of mammals and especially domestic dogs (e.g. Patronek et al., '97; Michell, '99; Egenvall et al., 2000; Speakman et al., 2003), longevity decreases with body size.

We explore two datasets for dogs and find support for a negative relationship between size and longevity if we consider variation across breeds. Within breeds, however, the relationship is not negative and is slightly, but significantly, positive in the larger of the two datasets. The negative across-breed relationship is probably the consequence of short life spans in large breeds. Artificial selection for extremely high growth rates in large breeds appears to have led to developmental diseases that seriously diminish longevity. *J. Exp. Zool. (Mol. Dev. Evol.)* 308B:119–126, 2007. © 2006 Wiley-Liss, Inc.

How to cite this article: Galis F, Sluijs IVD, Dooren TJMV, Metz JAJ, Nussbaumer M. 2007. Do large dogs die young? *J. Exp. Zool. (Mol. Dev. Evol.)* 308B:119–126.

The commonly found positive interspecific relationship between size and longevity can be explained relatively well with the oxidative stress theory of aging (Harman, '56; Beckman and Ames, '98). This theory postulates that aging is linked with energy expenditure because of cellular damage induced by free radicals that are a by-product of oxidative metabolism. Speakman et al. (2002) have indeed found a negative interspecific relationship between energy use and longevity in mammals. Since small mammalian species in general have a higher mass-specific metabolic rate than large species, a positive interspecific relationship between size and longevity would be expected. Within species small adult individuals also have higher metabolic rates than large individuals (Burger and Johnson, '91; Speakman et al., 2003). This fact taken by itself leads to an expectation of a positive intraspecific relationship between size and longevity (Speakman et al., 2003).

Several other hypotheses have been proposed to explain why some species live longer than others, given their size and metabolic rate. The “mutation accumulation” theory of Medawar ('52) proposes that populations that experience high mortality

rates accumulate deleterious mutations that reduce fitness late in life, because purifying selection has little power to remove late-acting mutations from the gene pool. The “antagonistic pleiotropy” hypothesis of Williams ('57) proposes that high mortality rates will select for earlier maturity and a higher rate of investment in reproduction early in life, even if this incurs a cost later in life. The “disposable soma” hypothesis of Kirkwood ('77) assumes that anti-aging mechanisms are costly and that, therefore, selection for anti-aging mechanisms will vary depending on the strength of extrinsic mortality. When extrinsic mortality is high and animals invariably die young, anti-aging mechanisms such as lower free radical production and better avoidance and repair mechanisms will have little impact on life span and thus will not be favored. However, when extrinsic mortality is low,

Grant sponsor: Dutch NWO-VENI grant.

*Correspondence to: F. Galis, Institute of Biology, Leiden University, P.O. Box 9516, 2300 RA Leiden, The Netherlands.
E-mail: galis@rulsfb.leidenuniv.nl

Received 1 March 2006; Accepted 18 April 2006

Published online 20 June 2006 in Wiley InterScience (www.interscience.wiley.com). DOI: 10.1002/jez.b. 21116

anti-aging mechanisms may have a substantial impact on life span and, therefore, a strong selective advantage. Although practical limitations often constrain the choice of species for comparative gerontological analyses (Rose, '91; Speakman et al., 2002), considerable support for the latter hypothesis has now accumulated (Austad, '93; Ku et al., '93; Barja et al., '94; Cortopassi and Wang, '96; Ogburn et al., '98; Kapahi et al., '99; Ricklefs and Scheuerlein, 2001; Blanco and Sherman, 2005). Recently, it has been shown for guppies that the relationship between extrinsic mortality and longevity may be more complex, with strong predation leading to a high rate of aging late in life, but to a low rate of aging earlier on (Reznick et al., 2004). Yet another hypothesis does not concern extrinsic mortality rates but proposes that a high growth rate will shorten life spans by increasing free-radical production (reviewed in Rollo, 2002). Growth rate indeed appears to be negatively associated with longevity (Ricklefs, '93; Olsson and Shine, 2002; Reznick et al., 2002; Rollo, 2002; Metcalfe and Monaghan, 2003, but see Anisimov, 2004).

Several authors (Austad, '97; Rollo, 2002; Speakman et al., 2003) have concluded that size and longevity may be negatively correlated within species of mammals. Support comes from rodents in which small size was induced by a calorie-restricted diet or by mutations resulting in a low growth rate (Rollo, 2002). In nature, however, a large size might also be the result of a protracted growth period rather than of a fast growth rate.

Most other support comes from studies on dogs. In dogs, small individuals have a much higher mass-specific metabolic rate than large ones (Burger and Johnson, '91; Speakman et al., 2003). There is no indication that small dogs have been selected for anti-aging mechanisms that could explain their longer life spans. A negative intraspecific relationship, therefore, provides a challenge for the oxidative stress theory of aging (Speakman et al., 2003), unless high growth rates in large dogs would explain the shorter life spans (Rollo, 2002). It is not known however, to what extent the differences in life span between small and large dogs may be confounded by genetic differences between small and large breeds. Strong selection and inbreeding have led to genetic differences between breeds (e.g., Ubbink et al., '98). To evaluate the influence of the differences between breeds, we have investigated the relationship between adult size and longevity across breeds and within breeds in two datasets.

One dataset (Veterinary Medical DataBase, VMDB) recorded weight (in classes) as a size measure and the other (Natural History Museum Bern, NMBE) a precise length measure in the skull that is highly correlated with other skeletal length measurements (Lüps, '74).

MATERIAL AND METHODS

Datasets

Data on size and longevity were obtained from the VMDB and the NMBE. We used longevity and weight measurements from 44,363 dogs from 134 breeds at the VMDB (longevity and weight (at death) measures in categories, longevity (years): 1–2, 2–4, 4–7, 7–10, 10–15, 15+; weight (kg): 0–0.5, 0.5–2.3, 2.3–6.8, 6.8–13.6, 13.6–22.7, 22.7–34.0, 34.0–45.4, 45.4+). We analysed those by using midpoint values of each category, except for the uppermost categories, where we used the lower bound of that category, because no upper limit was given. We only included breeds with individuals in at least three weight classes. The NMBE dataset consists of precise data on 859 dogs from 42 breeds. The length of the base of the brain stem (in mm) was taken as a measure of size for the dogs in the NMBE collection. This measure correlates highly with the length of the vertebral column, femur, pelvis and skull in most breeds (Lüps, '74). Breeds with a low correlation between the length of the base of the brain stem and other length measures were excluded from the dataset (Chihuahua, Greyhound, Bulldog, Boxer, Chow Chow, Bullterrier, Borzoi, French Bulldog, Akita, Pug, Dachshund, see Lüps, '74).

Age at death is recorded in months. Dogs that were known to have died in an accident, euthanized for behavioural problems or that were younger than 1 year old were not included in the dataset.

Statistical analysis

The data were analysed using bivariate linear random effect models (Meyer, '85). For both datasets, the same procedure was followed. Per trait y and per sex, we estimated parameters of a model of the form $y_{ij} = \mu + z_i + e_j$ with μ is the mean of that data subset, z a random effect specific to the i th breed and e the residual error within breeds (indexed by individuals j). Between sexes and for the same trait, the random breed effects were assumed to be the same. When we investi-

gated whether that assumption was warranted using single-trait analysis, we found that sex-specific random effects were not significantly different between sexes.

In the bivariate analysis, we estimated a variance–covariance matrix of the breed effects for life span and size (weight or length), and a variance–covariance matrix of the residual within-breed error terms. Estimation was performed using ASReml software for mixed linear models (Gilmour et al., 2002). Standard errors were calculated from the estimated Fisher information matrix. We tested for significant differences from zero for the variance components using *t*-tests (Table 1, Coltman et al., 2001). Two-sided *P*-values are reported. We also did likelihood ratio tests for significance of the covariances in the bivariate model, and conservative likelihood ratio tests for the breed variances in univariate models (Pinheiro and Bates, 2000), which are in agreement with the *t*-tests.

The VMDB dataset has a relatively low number of life span and weight classes as variables. Therefore the measurement error is large. In addition, both emaciated and obese dogs will

influence the relationship. However, we believe that the very large size of the dataset makes the conclusions we draw reliable. We treated breed effects as independent and did not correct for phylogenetic correlations, because of the highly reticulate nature of the evolution of most dog breeds (see Discussion).

RELATIONSHIP BETWEEN SIZE AND LONGEVITY

We find negative correlations between life span and size for variation between breeds (see Fig. 1 and Table 1), but overall positive correlations within breeds (see Table 1, see also Fig. 2). In other words, females and males of larger and heavier breeds die younger, but within breeds larger and heavier individuals die older on average, with the proviso that the correlations are only significant for the large VMDB dataset and that the correlations within breeds are much lower than between breeds. Figure 1 suggests that, in the NMBE dataset, size might have a non-linear relationship with life span, since very small dogs seem to have reduced life span too. However,




Fig. 1. Dogs from large breeds usually die younger than dogs from small breeds. Life span and size effects are shown per breed, i.e., the predicted life span of a breed relative to the overall mean, corrected for sex (as predicted by a random effects model, see Methods). Size is represented by bodyweight (kg) in the VMDB dataset (left) and by the length of the base of the brain stem (mm) in the NMBE dataset (right), life span is in years. The area of each point is proportional to the sample size per breed. Maximum sample size per breed is 97 in the NMBE dataset, 3,378 in the VMDB dataset. Minimum sample sizes are 4 (NMBE) and 5 (VMDB).

TABLE 1. Variances of lifespan and size (weight or length) between and within breeds

| VMDB | Lifespan | Weight |
|------------------------------------|---------------------------------|-----------------------------------|
| <i>Between breeds</i> | | |
| Covariance matrix breed effects | | |
| Lifespan | 1.22 (s.e. 0.18, $P < 0.001$) | -6.47 (s.e. 1.26, $P < 0.001$) |
| Weight | $r = -0.54$ | 116.20 (s.e. 14.41, $P < 0.001$) |
| <i>Within breeds</i> | | |
| Covariance matrix residual effects | | |
| Lifespan | 14.15 (s.e. 0.01, $P < 0.001$) | 1.73 (s.e. 0.13, $P < 0.001$) |
| Weight | $r = 0.06$ | 50.32 (s.e. 0.34, $P < 0.001$) |
| NMBE | Lifespan | Length |
| <i>Between breeds</i> | | |
| Covariance matrix breed effects | | |
| Lifespan | 3.08 (s.e. 0.95, $P = 0.002$) | -6.85 (s.e. 3.74, $P = 0.07$) |
| Length | $r = -0.36$ | 116.20 (s.e. 25.93, $P < 0.001$) |
| <i>Within breeds</i> | | |
| Covariance matrix residual effects | | |
| Lifespan | 12.79 (s.e. 0.72, $P < 0.001$) | 0.57 (s.e. 0.45, $P = 0.21$) |
| Length | $r = 0.05$ | 10.05 (s.e. 0.57, $P < 0.001$) |




Fig. 2. Funnel graphs of Pearson correlation coefficients between life span and size within breeds are shown (solid circles: males; open circles: females). Size is represented by body weight in kg in the VMDB dataset (left) and by the length of the base of the brain stem in mm in the NMBE dataset (right), life span is in years. Correlations are plotted as a function of sample size per breed. A horizontal line is drawn at correlation zero, and the two dotted lines of the funnel correspond to the threshold where negative and positive correlation estimates become significantly different from zero for a specific sample size. Sample correlations outside the funnel are significantly different from zero. In the VMDB dataset, there is a clear tendency towards positive correlations, corresponding to the significant test in Table 1. This trend is most clearly visible at large sample sizes. We found five significantly negative and 49 positive correlation coefficients (3 vs. 27 in males; 2 vs. 22 in females) in the VMDB dataset, two significantly positive estimates in the NMBE dataset (both in males). Overall, correlation estimates are 41 times positive and 21 negative in the NMBE dataset and 185 times positive and 70 negative in the VMDB dataset.

the same breeds are represented in the VMDB dataset and no reduced life span is visible for very small dogs in this dataset.

DISCUSSION

No negative relationship within breeds: We found a slightly positive relationship within breeds between size and longevity in our datasets (see Table 1), but the relationship is only significant in the larger dataset (VMDB). Similarly, the negative association across breeds is only significant in the larger dataset (negative trend for the NMBE). The discrepancy between the two datasets is most probably due to the difference in sample size (Fig. 2). The estimated correlation coefficients have similar values. Within breeds large dogs do not die younger than small ones, contrary to the assumption in the literature. Other data on within-breed and within-strain comparisons show no significant relationship (Patronek et al., '97; Speakman et al., 2002). However, Miller et al. (2002) found a negative relationship between size and longevity in a population of lab mice. This population, though, was composed of four different inbred mouse strains and the results may,

therefore, have been confounded by genetic differences between strains (see also Anisimov et al., 2004 and Khazaeli et al., 2005 on the importance of differences between strains).

Phylogenetic angle: We did not correct our results for phylogenetic correlations, because the most complete and recent phylogenetic analysis does not reveal significant genetic differences between 78 of 85 breeds (Parker et al., 2004). This is presumably because the bifurcating tree model of the analysis is not a good approximation for the intensely reticulate nature of the evolution of most dog breeds (Parker et al., 2004; Bannasch et al., 2005; see also Vilà et al., 2005). A particularly striking example is provided by the Irish Wolfhound which is supposedly a mix of Glengarry Deerhounds, Borzois, Great Danes, Tibetan Mastiffs and perhaps also of original Irish Wolfhounds and some other breeds. Freckleton et al. (2002) conclude that the contribution of the phylogenetic signal tends to be small in such circumstances and may even be misleading. Finally, a check on separate Pearson correlation coefficients within individual breeds (Fig. 2) confirms our conclusion that, overall, larger dogs within breeds do not die younger than smaller




Fig. 3. Selection for large size during the last century has been successful in Saint Bernard Dogs, similarly to that in other large breeds. Top left: male, 1968; top right: female, 2001. Bottom left: male, 1893; bottom right: female, ca. 1880–1890.

dogs. There were few significantly negative correlations between size and longevity in the two datasets and many more significantly positive correlation coefficients. The amount of significant negatives, five, is close to the average amount of type I errors expected, such that these results reflect the slightly positive trend of our mixed model analysis (Table 1). For most breeds there was no significant relationship and in both datasets there were significantly more positive correlation coefficients than negative ones, reflecting the slightly positive trend of our analysis (Fig. 2).

Why do dogs from large breeds die young? Dogs from large breeds usually die around the age of 6 years, which is young for dogs in general (and for wolves, Mech, '70; MacDonald, '84). This early mortality cannot be explained by oxidative damage due to size-related energy expenditure because dogs from large breeds have a lower mass-specific metabolic rate than dogs from small breeds (Burger and Johnson, '91; Speakman et al., 2003). In addition, there is no indication that breeds were selected for anti-aging mechanisms that could explain differences in mortality between breeds. Rollo (2002) has suggested that the elevated mortality of large individuals might be caused by high growth rates, which would induce high rates of oxidative damage during early life. Indeed, growth rates in large breeds during the first year are very high. Great Danes increase in weight 100-fold from birth in the first year, compared to 60-fold in wolves in captivity, 20-fold in poodles and 3-fold in humans (Mech, '70; Hawthorne et al., 2004). The proposal that a high free-radical production is involved in the early mortality is in agreement with extremely high rates of bone cancer in large breeds, 60–100-fold that of smaller breeds (Tjalma, '66; Withrow et al., '91). In addition, the high plasma levels of the growth-promoting insulin-like growth factor I (Igf-1) that are found in large breeds (Eigenmann et al., '88; Tryfonidou et al., 2003), combined with the inverse relation between Insulin/Igf-1 signaling and longevity in invertebrates and probably vertebrates (Partridge and Gems, 2002; Barbieri et al., 2003; Holzenberger et al., 2003; but see Carter et al., 2002) supports the idea that high growth rates cause the early mortality in large dog breeds.

However, when deaths from free-radical-associated diseases such as cancer and cardiovascular diseases are excluded, the average age at death of giant breeds is not increased, at least for Irish

Wolfhounds and St. Bernard Dogs (Bernardi, '88; SBCA Survey, '93). The oxidative stress theory of aging can, thus, only in part explain the early mortality. Additional important factors in the early death of dogs from such large breeds are developmental skeletal diseases, such as hip dysplasia and osteochondrosis (Dämmrich, '91; Slater et al., '92; Kealy et al., 2002). These diseases are also linked to high growth rates and appear to be due to a mismatch between the rate of weight increase and skeletal development and growth. The situation in large breeds is so unnatural that drinking ad libitum from the mother leads to a considerably increased incidence of joint diseases, when compared to a reduced intake of milk from bottles (Slater et al., '92). The high growth rates are presumably the result of artificial selection, as a side effect of selection for large mature size (Dämmrich, '91). In this respect, it is of interest to note that in *Drosophila* extreme artificial selection for rapid development has also led to pathological conditions and early mortality (Chippindale et al., '97). The size of giant dog breeds (Great Dane, Newfoundland, St. Bernard dog, Irish Wolfhound) has remarkably increased since 1800–1900 (see Fig. 3). For instance, the breed standard for St. Bernard dogs now specifies a shoulder height of between 70 and 90 cm and these dogs weigh 65–85 kg, whereas a typical 19th century dog was approximately 60 cm high and weighed less than 50 kg (Nussbaumer, 2000). The negative traits associated with the high growth rates would, presumably, be strongly selected against in nature. Only the relaxed selection due to human care allows these traits to persist.

The early mortality in large dog breeds, thus, does not appear to pose a threat to the oxidative stress theory of aging. Artificial selection on size has apparently led to pathological conditions in large breeds that misleadingly suggests that large body sizes negatively affect life span in dogs. Our study shows that research on aging and other fitness-related parameters may easily be flawed if no attention is given to the confounding effects of differences in the genetic backgrounds of breeds and strains (see also Anisimov et al., 2004; Khazaeli et al., 2005). This is particularly relevant because artificial selection has played such an important role in the species that are most often used for experimentation. Hence, for a better understanding of the intraspecific relationship between size and longevity in mammals, studies on natural populations are eagerly awaited.

ACKNOWLEDGMENTS

We thank Jacques van Alphen, Ricardo Azevedo, Russ Lande and especially Ray Huey for discussions and comments, Marge Macgregor for making available to us the health survey from the Saint Bernard Club of America and Rudolf Thomann from the Swiss Saint Bernard Dog Club for information on current weights of St. Bernard Dogs and the Veterinary Medical DataBase (VMDB) for making available their dataset. TVD was supported by a Dutch NWO-VENI grant.

LITERATURE CITED

- Anisimov VN, Arbeev KG, Popovich IG, Zabezhinski MA, Rosenfeld SV, Piskunova TS, Arbeeva LS, Semenchenko AV, Yashin AI. 2004. Body weight is not always a good predictor of longevity in mice. *Exp Gerontol* 39:305–319.
- Austad SN. 1993. Retarded senescence in an insular population of Virginia opossums (*Didelphis virginiana*). *J Zool* 229:695–708.
- Austad SN. 1997. Comparative aging and life histories in mammals. *Exp Gerontol* 32:23–38.
- Bannasch DL, Bannasch MJ, Ryun JR, Famula TR, Pedersen NC. 2005. Y chromosome haplotypes analysis in purebred dogs. *Mammalian Genome* 16:273–280.
- Barbieri M, Bonafe M, Franceschi C, Paolisso G. 2003. Insulin/IGF-I-signaling pathway: an evolutionarily conserved mechanism of longevity from yeast to humans. *Am J Physiol Endocrinol Metab* 285:E1064–E1071.
- Barja G, Cadenas S, Rojas C, Perez-Campo R, Lopez-Torres M. 1994. Low mitochondrial free radical production per unit O₂ consumption can explain the simultaneous presence of high longevity and high aerobic metabolic rate in birds. *Free Radical Res* 21:317–328.
- Beckman KB, Ames BN. 1998. The free radical theory of aging matures. *Physiol Rev* 78:547–581.
- Bernardi G. 1988. Longevity and mortality in the Irish Wolfhound in the United States: 1966–1986. *Harp Hound* 78–84.
- Blanco MA, Sherman PW. 2005. Maximum longevity of chemically protected and non-protected fishes, reptiles, and amphibians support evolutionary hypotheses of aging. *Mech Ageing Dev* 126:794–803.
- Burger IH, Johnson JV. 1991. Dogs large and small: the allometry of energy requirements within a single species. *J Nutr* 121 (11 suppl):S18–S21.
- Carter CS, Ramsey MM, Sonntag WE. 2002. A critical analysis of the role of growth hormone and IGF-1 in aging and lifespan. *Trends Genet* 18:295–301.
- Chippendale AK, Alipaz JA, Chen HW, Rose MR. 1997. Experimental evolution of accelerated development in *Drosophila*. Developmental speed and larval survival. *Evolution* 51:1536–1551.
- Coltman DW, Pilkington J, Kruuk LEB, Wilson K, Pemberton JM. 2001. Positive genetic correlation between parasite resistance and body size in a free-living ungulate population. *Evolution* 55:2116–2125.
- Cortopassi GA, Wang E. 1996. There is substantial agreement among interspecies estimates of DNA repair activity. *Mech Ageing Dev* 91:211–218.
- Dämmrich K. 1991. Relationship between nutrition and bone growth in large and giant dogs. *J Nutr* 121:433–446.
- Egenvall A, Bonnett BN, Shoukri M, Olson P, Hedhammer Å, Dohoo I. 2000. Age pattern of mortality in eight breeds of insured dogs in Sweden. *Prev Vet Med* 46:1–14.
- Eigenmann JE, Amador A, Patterson DF. 1988. Insulin-like growth factor I levels in proportionate dogs, chondro-dystrophic dogs and in giant dogs. *Acta Endocrinol* 118:105–108.
- Freckleton RP, Harvey PH, Pagel M. 2002. Phylogenetic analysis and comparative data: a test and review of evidence. *Am Nat* 160:712–726.
- Gilmour AR, Cullis BR, Welham SJ, Thompson R. 2002. ASReml reference manual, 2nd edition, Release 1.0 NSW Agriculture Biometrical Bulletin 3, NSW Agriculture, Locked Bag, Orange, NSW 2800, Australia.
- Harman D. 1956. Aging, a theory based on free radical and radiation chemistry. *J Gerontol* 11:298–300.
- Hawthorne AJ, Booles D, Nugent PA, Gettinby G, Wilkinson J. 2004. Body-weight changes during growth in puppies of different breeds. *J Nutr* 134:2027S–2030S.
- Holzenberger M, Dupont J, Ducos B, Leneuve P, Geloën A, Evens P, Cervera P, Le Bouc Y. 2003. IGF-1 receptor regulates lifespan and resistance to oxidative stress in mice. *Nature* 421:182–187.
- Kapahi P, Boulton ME, Kirkwood TB. 1999. Positive correlation between mammalian life span and cellular resistance to stress. *Free Radical Biol Med* 26:495–500.
- Kealy RD, Lawler DF, Ballam JM, Mantz SL, Biery DN, Greeley EH, Lust G, Segre M, Smith GK, Stowe HD. 2002. Effects of diet restriction on life span and age-related changes in dogs. *J Am Vet Med Assoc* 220:1315–1320.
- Khazaeli AA, Van Voorhies W, Curtisinger JW. 2005. The relationship between lifespan and adult body size is highly strain-specific in *Drosophila melanogaster*. *Exp Gerontol* 40:377–385.
- Kirkwood TBL. 1977. Evolution of aging. *Nature* 270:301–304.
- Ku H-H, Brunk UT, Sohal RS. 1993. Relationship between mitochondrial superoxide and hydrogen peroxide production and longevity of mammalian species. *Free Radical Biol Med* 15:621–627.
- Lüps P. 1974. Biometrische Untersuchungen an der Schädelbasis des Haushundes. *Zool Anz* 5/6:383–413.
- MacDonald D. 1984. The encyclopedia of mammals, MacDonald, D., editor series. New York.
- Mech LD. 1970. The wolf. The ecology and behavior of an endangered species. Minneapolis: University of Minnesota Press.
- Medawar PB. 1952. An unsolved problem of biology. London: Lewis.
- Metcalfe NB, Monaghan P. 2003. Growth versus lifespan: perspectives from evolutionary ecology. *Exp Gerontol* 38:935–940.
- Meyer K. 1985. Maximum likelihood estimation of variance components for a multivariate mixed model with equal design matrices. *Biometrics* 41:153–166.
- Michell AR. 1999. Longevity of British breeds of dog and its relationship with sex, size, cardiovascular variables and disease. *Vet Rec* 145:625–629.
- Miller RA, Harper JM, Galecki A, Burke DT. 2002. Big mice die young: early life body weight predicts longevity in genetically heterogeneous mice. *Aging Cell* 1:22–29.

- Nussbaumer M. 2000. Barry vom Grossen St. Bernard. Bern, Switzerland: Naturhistorisches Museum der Burgergemeinde.
- Ogburn CE, Austad SN, Holmes DJ, Kiklevich JV, Gollahon K, Rabonovitch PS, Martin GM. 1998. Cultured renal epithelial cells from birds and mice: enhanced resistance of avian cells to oxidative stress and DNA damage. *J Gerontol* 53:B287–B292.
- Olsson M, Shine R. 2002. Growth to death in lizards. *Evolution* 56:1867–1867.
- Parker HG, Kim LV, Sutter NB, Carlson S, Llorentzen TD, Malek TB, Johnson GS, DeFrance HB, Ostrander EA, Kruglyak L. 2004. Genetic structure of the purebred domestic dog. *Science* 304:1160–1164.
- Partridge L, Gems D. 2002. Mechanisms of ageing: public or private? *Nat Rev* 3:165–175.
- Patronek GJ, Waters DJ, Glickman LT. 1997. Comparative longevity of pet dogs and humans: Implications for gerontology research. *J Gerontol Biol Sci* 52A:B171–B178.
- Pinheiro J, Bates DM. 2000. Mixed effects models in S and S-plus. New York: Springer-Verlag.
- Reznick DN, Ghalambor CK, Nunney L. 2002. The evolution of senescence in fish. *Mech Aging Dev* 123:773–789.
- Reznick DN, Bryant MJ, Roff D, Ghalambor CK, Ghalambor DE. 2004. Effect of extrinsic mortality on the evolution of senescence in guppies. *Nature* 431:1095–1099.
- Ricklefs RE. 1993. Sibling competition, hatching asynchrony, incubation period, and lifespan in altricial birds. *Curr Ornithol* 11:199–276.
- Ricklefs RE, Scheuerlein A. 2001. Comparison of aging-related mortality among birds and mammals. *Exp Gerontol* 36:845–857.
- Rollo CD. 2002. Growth negatively impacts the life span of animals. *Evol Dev* 4:55–61.
- Rose MR. 1991. Evolutionary biology of aging. Oxford: Oxford University Press.
- Saint Bernard Club of America. 1993. SBCA health survey. *Saint Fancier March/April*:59–73.
- Slater MR, Scarlett JM, Donohuey S, Kaderly RE, Bonnett BN, Cockshutt J, Erb HN. 1992. Diet and exercise as potential risk factors for osteochondritis dissecans in dogs. *Am J Vet Res* 53:2119–2124.
- Speakman JR, Selman C, McLaren JS, Harper EJ. 2002. Living fast, dying when? The link between aging and energetics. *J Nutr* 132(6, Suppl 2):1583S–1597S.
- Speakman JR, van Acker A, Harper EJ. 2003. Age-related changes in the metabolism and body composition of three dog breeds and their relationship to life expectancy. *Aging Cell* 2:265–275.
- Tjalma RA. 1966. Canine bone sarcoma: estimation of relative risk as function of body size. *J Natl Cancer Inst* 36:1137–1150.
- Tryfonidou MA, Holl MS, Vastenburg M, Oosterlaken-Dijksterhuis MA, Birkenhager-Frenkel DH, van den Brom WE, Hazewinkel HA. 2003. Hormonal regulation of calcium homeostasis in two breeds of dogs during growth at different rates. *J Anim Sci* 81:1568–1580.
- Ubbink GJ, van de Broek J, Hazewinkel HA, Rothuizen J. 1998. Cluster analysis of the genetic heterogeneity and disease distributions in purebred dog populations. *Vet Rec* 142:209–213.
- Vilà C, Seddon J, Ellegren H. 2005. Genes of domestic mammals augmented by backcrossing with wild ancestors. *Trends Genet* 21:214–218.
- Williams GC. 1957. Pleiotropy, natural selection and the evolution of senescence. *Evolution* 11:398–411.
- Withrow SJ, Powers BE, Straw RC, Wilkins RM. 1991. Comparative aspects of osteosarcoma: dog versus man. *Clin Orthop* 270:159–168.